



Walton, E., Hibar, D. P., van Erp, T. G. M., Potkin, S. G., Roiz-Santiañez, R., Crespo-Facorro, B., Suarez-Pinilla, P., van Haren, N. E. M., de Zwarte, S. M. C., Kahn, R. S., Cahn, W., Doan, N. T., Jørgensen, K. N., Gurholt, T. P., Agartz, I., Andreassen, O. A., Westlye, L. T., Melle, I., Berg, A. O. (2018). Prefrontal cortical thinning links to negative symptoms in schizophrenia via the ENIGMA consortium. *Psychological Medicine*, 48(1), 82-94.
<https://doi.org/10.1017/S0033291717001283>

Peer reviewed version

Link to published version (if available):
[10.1017/S0033291717001283](https://doi.org/10.1017/S0033291717001283)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cambridge Core at <https://www.cambridge.org/core/journals/psychological-medicine/article/prefrontal-cortical-thinning-links-to-negative-symptoms-in-schizophrenia-via-the-enigma-consortium/B7FE158F6DEAE030607B51B586C5C553> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Self-archiving version

Prefrontal cortical thinning links to negative symptoms in schizophrenia via the ENIGMA consortium

Esther Walton^{1,2,3}; Derrek P Hibar⁴; Theo GM van Erp⁵; Steven G Potkin⁵; Roberto Roiz-Santiañez^{6,7}; Benedicto Crespo-Facorro^{6,7}; Paula Suarez-Pinilla^{6,7}; Neeltje EM Van Haren⁸; Sonja MC de Zwarte⁸; Rene S Kahn⁸; Wiepke Cahn⁸; Nhat Trung Doan⁹; Kjetil N Jørgensen^{9,10}; Tiril P Gurholt⁹; Ingrid Agartz^{9,10,11}; Ole A Andreassen^{12,9}; Lars T Westlye¹²; Ingrid Melle^{9,12}; Akiah O Berg⁹; Lynn Morch-Johnsen^{9,10}; Ann Færden¹³; Lena Flyckt¹⁴; Helena Fatouros-Bergman¹⁴; Karolinska Schizophrenia Project Consortium (KaSP)¹⁵; Erik G Jönsson^{9,11}; Ryota Hashimoto^{16,17}; Hidenaga Yamamori¹⁷; Masaki Fukunaga¹⁸; Neda Jahanshad⁴; Pietro De Rossi^{19,20}; Fabrizio Piras²⁰; Nerisa Banaj²⁰; Gianfranco Spalletta^{20,21}; Raquel E Gur²²; Ruben C Gur²²; Daniel H Wolf²²; Theodore D Satterthwaite²²; Lauren M Beard²²; Iris E Sommer⁸; Sanne Koops⁸; Oliver Gruber²³; Anja Richter²³; Bernd Krämer²³; Sinead Kelly^{4,24}; Gary Donohoe²⁵; Colm McDonald²⁵; Dara M Cannon²⁵; Aiden Corvin²⁴; Michael Gill²⁴; Annabella Di Giorgio²⁶; Alessandro Bertolino²⁷; Stephen Lawrie²⁸; Thomas Nickson²⁸; Heather C Whalley²⁸; Emma Neilson²⁸; Vince D Calhoun^{29,30}; Paul M Thompson⁴; Jessica A Turner³¹; Stefan Ehrlich²

1 Department of Psychology, Georgia State University, Atlanta GA 30302

2 Division of Psychological and Social Medicine and Developmental Neurosciences, Faculty of Medicine, Technische Universität Dresden, Fetscherstr. 74, 01307 Dresden, Germany

3 Department of Psychology, Institute of Psychology, Psychiatry and Neuroscience, King's College London, London, SE5 8AF, United Kingdom

4 Imaging Genetics Center, Keck School of Medicine, University of Southern California, Marina del Rey, CA, United States

5 Department of Psychiatry and Human Behavior, University of California, Irvine, California

6 Department of Psychiatry, University Hospital Marqués de Valdecilla, School of Medicine, University of Cantabria–IDIVAL, Avda. Valdecilla s/n, 39008, Santander, Spain

7 Cibersam (Centro Investigación Biomédica en Red Salud Mental), Avda. Valdecilla s/n, 39008, Santander, Spain

8 Department of Psychiatry, Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht, The Netherlands

9 NORMENT, KG Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, University of Oslo, P.O. Box 4956 Nydalen, 0424 Oslo, Norway

10 Department of Psychiatric Research, Diakonhjemmet Hospital, P.O. Box 85 Vinderen,
0319 Oslo, Norway

11 Department of Clinical Neuroscience, Karolinska Institutet, 171 77 Stockholm, Sweden

12 NORMENT, KG Jebsen Centre for Psychosis Research, Division of Mental Health and
Addiction, Oslo University Hospital, P.O. Box 4956 Nydalen, 0424, Oslo, Norway

13 Division of Mental Health and Addiction, Oslo University Hospital, P.O. Box 4956 Nydalen,
0424, Oslo, Norway

14 Karolinska Institutet, Department of Clinical Neuroscience, Centre for Psychiatry
Research, Norra Stationsgatan 69, 113 64 Stockholm, Sweden

15 Members of Karolinska Schizophrenia Project (KaSP) are listed at the end of the article as
collaborators

16 Molecular Research Center for Children's Mental Development, United Graduate School
of Child Development, Osaka University D3, 2-2, Yamadaoka, Suita, Osaka, 565-0871, Japan

17 Department of Psychiatry, Osaka University Graduate School of Medicine D3, 2-2,
Yamadaoka, Suita, Osaka, 565-0871, Japan

18 Division of Cerebral Integration, National Institute for Physiological Sciences, 38
Nishigonaka Myodaiji, Okazaki, Aichi, 444-8585, Japan

19 NESMOS Department (Neurosciences, Mental Health and Sensory Functions), School of Medicine and Psychology, Sapienza University, Rome, Italy

20 Department of Clinical and Behavioural Neurology, IRCCS Santa Lucia Foundation, 00179, Rome, Italy

21 Beth K. and Stuart C. Yudofsky Division of Neuropsychiatry Menninger Department of Psychiatry and Behavioral Sciences Baylor College of Medicine Houston, TX, USA

22 Brain Behavior Laboratory, University of Pennsylvania, Philadelphia PA USA 19104

23 Center for Translational Research in Systems Neuroscience and Psychiatry, Department of Psychiatry and Psychotherapy, University Medical Center Göttingen, Von-Siebold-Str. 5, 37075 Göttingen, Germany

24 Trinity College, Dublin, Ireland

25 Neuroimaging and Cognitive Genomics Centre, NCBES Galway Neuroscience Centre, National University of Ireland Galway, Galway, Ireland

26 Section of Psychiatry and Psychology, IRCCS Casa Sollievo della Sofferenza, S.G. Rotondo (FG), 71013 Italy

27 Psychiatric Neuroscience Group, University of Bari 'Aldo Moro', Bari, 70124 Italy

28 Division of Psychiatry, University of Edinburgh, Royal Edinburgh Hospital, Morningside, Edinburgh, EH10 5HF

29 The Mind Research Network, Albuquerque, NM 87106, United States

30 Department of Electrical and Computer Engineering, University of New Mexico,
Albuquerque, NM 87131, United States

31 Department of Psychology and Neuroscience Institute, Georgia State University, Atlanta
GA 30302

Corresponding author:

Stefan Ehrlich, M.D; Division of Psychological and Social Medicine and Developmental
Neurosciences, Faculty of Medicine, Technische Universität Dresden, Fetscherstr. 74, 01307
Dresden, Germany, Phone: +49 (0)351 458-2244, Fax: +49 (0)351 458 -5754, Email:
stefan.ehrlich@tu-dresden.de

Word count

Abstract: 242

Manuscript: 3 722

Number of Tables: 1

Number of Figures: 1

Number of Supplementary Materials: 2

Keywords: schizophrenia, negative symptoms, medial orbitofrontal cortex, cortical thickness, ENIGMA, FreeSurfer, MRI, SANS, PANSS

Financial support

ENIGMA was supported in part by a Consortium grant (U54 EB020403 to PMT) from the NIH Institutes contributing to the Big Data to Knowledge (BD2K) Initiative, including the NIBIB and NCI. The authors would also like to express their gratitude for the personal support from the Deutsche Forschungsgemeinschaft (Research Fellowship to EW; Wa 3635/1-1). For additional support, see supplementary section 3.

Abstract

Background

Our understanding of the complex relationship between schizophrenia symptomatology and etiological factors can be improved by studying brain-based correlates of schizophrenia.

Research showed that impairments in value processing and executive functioning, which have been associated with prefrontal brain areas (particularly the medial orbitofrontal cortex (MOFC)), are linked to negative symptoms. Here we tested the hypothesis that MOFC thickness is associated with negative symptom severity.

Methods

This study included 1 985 individuals with schizophrenia from seventeen research groups around the world contributing to the ENIGMA Schizophrenia Working Group. Cortical thickness values were obtained from T1-weighted structural brain scans using FreeSurfer. A meta-analysis across sites was conducted over effect sizes from a model predicting cortical thickness by negative symptom score (harmonized SANS or PANSS scores).

Results

Meta-analytical results showed that left, but not right, MOFC thickness was significantly associated with negative symptom severity ($\beta_{\text{std}}=-0.075$; $p=0.019$) after accounting for age, gender and site. This effect remained significant ($p=0.036$) in a model including overall illness severity. Covarying for duration of illness, age of onset, antipsychotic medication or

handedness weakened the association of negative symptoms with left MOFC thickness. As part of a secondary analysis including ten other prefrontal regions further associations in the left lateral orbitofrontal gyrus and pars opercularis emerged.

Conclusions

Using an unusually large cohort and a meta-analytical approach, our findings point towards a link between prefrontal thinning and negative symptom severity in schizophrenia. This finding provides further insight into the relationship between structural brain abnormalities and negative symptoms in schizophrenia.

Introduction

Although advances have been made in our understanding of the pathophysiology of schizophrenia, the heterogeneity of the disorder impedes the effectiveness of biological and clinical research. The large number of cognitive and clinical symptoms within the syndrome and their considerable variability across patients likely reflects the impact of various etiological factors (Jablensky, 2006). Such variability makes it harder to achieve a comprehensive understanding underlying brain pathology. Investigations using large sample sizes to study symptoms associated with dimensions of behavior that can be measured quantitatively, may advance our understanding of brain behavior relationships within as well as across diagnostic categories. Investigating distinct symptoms of the disorder as a continuous variable may reveal more specific neurobiological mechanisms in schizophrenia.

Negative symptoms are characterized by flat or blunted affect, inability to experience pleasure (anhedonia), poverty of speech, lack of motivation and interest (avolition/apathy) and lack of desire to form relationships (Andreasen & Olsen, 1982). Patients with predominantly negative symptoms have poor pre-morbid adjustment during childhood or early adolescence and a low employment rate during adulthood, achieve low educational attainment and exhibit considerable cognitive impairment (Milev *et al.* 2005; Rosenheck *et al.* 2006; Jeppesen *et al.* 2008).

Studies have shown widespread cortical thickness reductions across the brain in patients with schizophrenia compared to healthy controls with frontal and temporal regions

being generally more affected than others areas (Nesvåg *et al.* 2008; Goldman, 2009; Schultz *et al.* 2010). In line with prior work, Ehrlich *et al.* (2012) and Geisler *et al.* (2015) recently reported marked reductions of cortical thickness in patients with schizophrenia, which were also related to executive functioning.

Executive functioning is often driven by incentives and motivation (Pessoa, 2009). When we make decisions between options with different (subjective) value or learn from positive or negative feedback, those processes are governed by a neural circuit that centers around structures such as the ventral striatum and the medial orbitofrontal cortex (MOFC) (Schlagenhauf *et al.* 2014; Deserno *et al.* 2016). Within this circuit, the MOFC is predominately linked to (subjective) value processing and positive affect (Burgdorf & Panksepp, 2006; Peters & Büchel, 2010; Grabenhorst & Rolls, 2011; Liu *et al.* 2011; Price & Harmon-Jones, 2011; Berridge & Kringelbach, 2015), which is impaired in schizophrenia (Cohen & Minor, 2010; Kalkstein *et al.* 2010). MOFC lesions in humans relate to deficits in reward-based learning, which can result in apathy and lack of affect (Hornak *et al.* 2003; Fellows & Farah, 2005; Fisher *et al.* 2011; Kühn & Gallinat, 2012). Accordingly, functional and structural imaging studies in healthy cohorts have reported associations between MOFC activity (and thickness) and characteristics that are comparable to the negative symptoms observed in schizophrenia (Harvey *et al.* 2007; Ducharme *et al.* 2014). For instance, functional connectivity between the orbitofrontal cortex and the dorsolateral prefrontal cortex was found to be dependent on levels of motivation (Szatkowska *et al.* 2008).

Given these findings one could hypothesize that altered brain structure and functioning in the orbitofrontal cortex may represent one of the potential mechanisms underlying negative symptoms in schizophrenia. This is motivated by the idea that i) neural representations of decision values may not be adequately generated and ii) reward feedback might not get completely transformed into motivational drive for goal-directed behaviour (Barch & Dowd, 2010; Deserno *et al.* 2013).

Several functional imaging studies have assessed the brain-based correlates of negative symptoms in schizophrenia. An early PET study found lower perfusion of several brain regions – including the MOFC – during hedonic judgments of positive and negative visual stimuli in patients with schizophrenia (Plailly *et al.* 2006). More recent fMRI studies in patients with schizophrenia found neural responses in the medial prefrontal cortex to be exaggerated upon omission of expected reward but blunted upon receipt of unexpected reward (Schlagenhauf *et al.* 2009). Simon *et al.* (2015) showed that neural responses in the ventral striatum during a reward anticipation paradigm were negatively associated with apathy – a core negative symptom; while no associations between ventral striatal responses and positive symptoms were observed. Importantly, they found lower connectivity between the ventral striatum and the MOFC in individuals with schizophrenia compared with controls. Furthermore, in one of the aforementioned studies focussing on receipt of unexpected reward in schizophrenia, medial frontal cortical activation predicted task-related motivation, which in turn predicted anhedonia severity (Segarra *et al.* 2016). In line

with this, another study could show that activity in the orbitofrontal cortex during hedonic processing was negatively correlated with anhedonia severity in people with schizophrenia (Harvey *et al.* 2010).

With respect to structural imaging, gray matter density of the orbitofrontal cortex has also been associated with self-fulfillment achievement motivation in healthy individuals (Takeuchi *et al.* 2014). In patients with schizophrenia, Venkatasubramanian *et al.* (2008) found that lower left MOFC thickness was associated with higher negative symptom severity. Similar findings have been reported by Nenadic *et al.* (2015), providing further evidence for the involvement of medial prefrontal regions in negative symptoms; though some negative findings also exist (Crespo-Facorro *et al.* 2011; Xiao *et al.* 2013).

Reasons for these inconsistencies may be two-fold. First, most studies conducted to date have had rather moderate sample sizes, investigating on average about a hundred patients. Second, moderating effects of confounders such as antipsychotic medication, illness severity, or duration of illness that may influence the association between thickness and negative symptoms have not been investigated extensively. Analyses of larger samples will increase power allowing to understand prior inconsistencies in findings as well as to study potential moderator effects.

Here we set out to investigate the structural correlates of negative symptoms in a large meta-analysis including almost 2 000 individuals with schizophrenia. Based on previous structural imaging findings, we hypothesised that lower MOFC thickness is associated with

negative symptom severity in schizophrenia. As part of an exploratory analysis we also aimed to understand relationships with cortical thickness in other prefrontal brain regions.

Methods

Study samples

The current study includes a total of 1 985 individuals with schizophrenia or schizophrenia spectrum diagnoses (see also SM section 2.1, subsection s) from seventeen research groups around the world as part of the ENIGMA Schizophrenia Working Group as described previously (van Erp *et al.* 2015). Schizophrenia diagnosis was based on the Diagnostic and Statistical Manual of Mental Disorders (DSM, editions III-R or IV) or the International Classification of Diseases (ICD, edition 10) criteria using either the Structured Clinical Interview for DSM Disorders (SCID), the Comprehensive Assessment of Symptoms and History (CASH), the Present State Examination (PSE), and/or a review of case files/medical records by trained clinicians. All individuals had negative symptom ratings and structural imaging data available. Mean sample size at each research site was 117 patients (range: 23-244). See supplementary Table 1s for more details.

Each study sample was collected with participants' written informed consent approved by local Institutional Review Boards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and

institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. No individual subject imaging or clinical data were shared among the ENIGMA institutions.

Negative symptom measures and score conversion

Negative symptom severity was assessed using the Scale for the Assessment of Negative Symptoms (SANS) (Andreasen, 1983) and the Positive and Negative Syndrome Scale (PANSS) (Kay *et al.* 1987). Negative symptom scores were calculated as follows:

1. Total SANS (Composite) score = sum of SANS items 1-7, 9-12, 14-16, 18-21, and 23-24;
2. Global SANS (Summary) score = sum of SANS items 8, 13, 17, 22, and 25 (which include affective flattening, alogia, avolition, anhedonia, and attention global rating scores, respectively); or
3. PANSS Negative = sum of PANSS items 8-14.

To harmonize scores, we decided to convert all negative scores (i.e. PANSS Negative and Total SANS Composite scores) to Global SANS (Summary) scores following recommendations by Andreasen (1983) and using the algorithms published in van Erp *et al.* (2014). For additional details see supplementary Section 1.1.

Image acquisition and processing

Based on the previous literature (Venkatasubramanian *et al.* 2008; Nenadic *et al.* 2015) we followed a region-of-interest (ROI) approach, focusing on the medial orbitofrontal cortex (MOFC), while ten other prefrontal regions were considered in additional exploratory analyses (see SM section 2.2). Left and right MOFC thickness values – based on the Desikan-Killiany atlas (Desikan *et al.* 2006) - were obtained using FreeSurfer (<http://surfer.nmr.mgh.harvard.edu>) from high-resolution T1-weighted structural brain scans. Details on study type (single site or multisite), scanner vendor/strength/sequence, acquisition parameters and FreeSurfer versions used are provided in supplementary Table 1s. For quality control, histograms of MOFC thickness values were generated and outliers were visually inspected by overlaying their parcellation on the subjects' anatomical images. Only parcellations judged to be accurate upon visual inspection were subjected to statistical analyses (see supplementary Table 1s for information on outlier removal and Figure 1s for left and right MOFC thickness descriptives by sample).

Statistical analyses

Within each sample, an association of negative symptoms with left and right MOFC thickness was analyzed using univariate linear regression analysis (R's linear model function *lm*) predicting mean MOFC thickness by global SANS score. The main analysis included age

and gender as covariates. In cases of multi-site studies (FBIRN, MCIC, UMCU and Osaka) binary dummy covariates were included in the model to account for $n-1$ sites. For samples where information was available, secondary models were run separately with each of the following covariates: 1) current antipsychotic medication (by medication group: atypical/typical/both/none; and by using chlorpromazine (CPZ) equivalents as described in Woods *et al.* (2003), available in a subsample of $n = 1\,178$), 2) duration of illness, 3) age of onset (defined as onset of symptoms), 4) illness severity (measured using PANSS Total score), and 5) handedness (right/left/ambidextrous). Additional exploratory analyses were carried out to investigate the link between negative symptom severity and cortical thickness in schizophrenia spectrum subtypes and in ten additional frontal brain regions. Results were ranked according to effect size. In addition, Bonferroni correction was applied to account for multiple testing (see SM section 2.2 for more details). Analyses of individual subject data were performed by the site that contributed the sample, using code created within the ENIGMA collaboration.

Meta-analyses

From each sample, standardized regression coefficients were extracted from the main and secondary models as a measure of effect size for the left and right MOFC using the *lm.beta* function in the *lm.beta* R package (Behrendt, 2014). A meta-analysis was conducted over these effect sizes using the *rma* function in the R package *metaphor* (Viechtbauer,

2010). We meta-analyzed the estimates across sites by weighting Fisher's r-to-z transformed effect size values by sample size in a random-effects model using the default REML estimator. The same procedure was used to investigate the effects of age, gender, illness severity and duration of illness on MOFC thickness. For analyses, in which both left and right MOFC were analyzed, the significance threshold was corrected for two tests ($p=0.05/2=0.025$). Descriptives are weighted by the sample size at each site using the weighted.mean procedure in R (R Development Core Team 2008).

Due to between-site differences in study characteristics such as antipsychotic medication, handedness and single vs multisite status (supplementary Table 1s), we used moderator analyses to investigate between-sample differences.

Results

Demographics

Mean age (weighted by sample-size) across patient samples was 34 years (range: 28-43). Mean patient samples were 68% male (range: 55-76%). The weighted mean duration of illness across the patient groups was 10 years (range: 1-20) and mean age of onset was 24 years (range: 19-29). For samples where current antipsychotic type and dose information was available, the weighted percentage of patients on first-generation (typical), second-generation antipsychotics (atypical), both typical and atypical or no antipsychotic

medication was 11%, 71%, 9% and 9%. Ninety percent of patients were right-handed (range: 68-95), while only 8% (range: 2-14) were left-handed and 2% (range: 0-25) were ambidextrous (Table 1 and supplementary Table 1s).

Meta-analysis

The weighted mean global SANS scores across the samples was 7.91 (range: 2.86-12.90). Weighted mean MOFC thickness was 2.46 mm (range: 2.26-2.71) in the left hemisphere and 2.42 mm (range: 2.17-2.61) in the right hemisphere. Meta-analytical results showed that global SANS scores were negatively associated with left MOFC thickness ($\beta_{\text{std}} = -0.075$; $p_{\text{SANS}} = 0.019$; Figure 1) after accounting for age, gender and number of sites (if applicable). Funnel plot inspection gave no indication of bias (Egger's $p = 0.193$; supplementary Figure 2s). However, effect sizes were found to be heterogeneous ($Q(16) = 26.399$; $p = 0.049$; $I^2 = 42.38\%$). We found only a trend effect (but in the same direction) of global SANS on right MOFC thickness ($\beta_{\text{std}} = -0.064$; $p_{\text{SANS}} = 0.055$). For detailed results, see supplementary Section 2.1, sections a and b).

Effects of covariates and moderator analyses

We carried on investigating both within-sample and moderating between-sample effects of age, gender, illness severity, duration of illness and age of onset as well as

antipsychotic medication, handedness, schizophrenia spectrum subtypes and multisite status based on samples, for which this information was available (supplementary Table 1s).

While a meta-analysis of within-sample effects indicated that left MOFC thickness decreased with age ($\beta_{\text{std}}=-0.237$; $p<0.0001$) and was lower in women ($\beta_{\text{std}}=-0.090$; $p<0.0001$), the main association of global SANS and left MOFC thickness remained significant ($\beta_{\text{std}}=-0.075$; $p_{\text{SANS}}=0.019$, see main model above and supplementary section 2.1, subsections a, c and d).

Overall illness severity (although correlated with negative symptoms; $R^2 = 59.5\%$) did not associate with left MOFC thickness ($p=0.555$) after accounting for age, gender and site (if applicable), while the global SANS effect in the same model remained significant ($\beta_{\text{std}}=-0.113$; $p_{\text{SANS}}=0.036$; see also supplementary section 2.1, subsections e and f).

Duration of illness correlated negatively with left MOFC thickness (Fisher's $z=-0.145$; $p<0.001$), but not with global SANS score ($p=0.190$). While duration of illness also correlated strongly with age (Fisher's $z=0.834$; $p<0.001$), it did not associate with left MOFC thickness above and beyond age in the lower regression models ($p=0.199$). However, additionally accounting for duration of illness within each sample also reduced the main effect of SANS on left MOFC thickness ($p_{\text{SANS}}=0.068$; see supplementary section 2.1, subsections g and h). Similar diminishing effects were observed when investigating the influence of age of onset (which was highly correlated with duration of illness; Fisher's $z=-0.658$; $p=0.015$), antipsychotic medication and handedness within each sample, which themselves did not

moderate the global SANS – left MOFC thickness relationship between samples ($p_{AO}=0.737$; $p_{med}=0.447$; $p_{hand}=0.580$; see supplementary section 2.1, section i, j and k). Investigating the effects of antipsychotic medication further, we found that mean left MOFC thickness estimates ($p=0.007$), but not mean global SANS estimates ($p=0.454$) differed by antipsychotic medication type (i.e. percentages of patients in each medication group in each sample, supplementary section 2.1, subsections l and m). Specifically, compared to unmedicated patients, left MOFC reductions were larger at sites with a larger percentage of patients treated with atypical ($\beta_{std}=-0.006$; $p=0.002$) or typical AP ($\beta_{std}=-0.007$; $p=0.003$). Proportion of patients, who were treated with both antipsychotic medication types, was not associated with MOFC thickness ($p=0.444$; supplementary section 2.1, subsection l). The reductions observed in patients treated with atypical (but not typical) antipsychotic medication remained significant in analyses that included gender, age, or duration of illness as covariates (supplementary section 2.1, subsection n and o). Current medication in CPZ units was not linked to left MOFC thickness ($p_{CPZ}=0.170$), while the main effect of negative symptom severity remained significant ($\beta_{std}=-0.095$; $p_{SANS}=0.040$; supplementary section 2.1, subsection q and r). Our main findings also remained stable ($\beta_{std}=-0.078$; $p_{SANS}=0.022$), when restricting the analyses to patients with DSM-IV schizophrenia subtypes or schizoaffective / -phreniform disorder only (supplementary section 2.1, subsection s).

Effect on other frontal regions

Exploratory analyses were used to analyse the associations between negative symptoms and ten other frontal brain regions (separately for the left and right hemisphere), controlling for sex, age and site (if applicable). Regions with effects similar in size to the left MOFC were found in the left pars opercularis ($\beta_{\text{std}}=-0.082$), the left and right lateral orbitofrontal gyrus (left: $\beta_{\text{std}}=-0.076$, right: $\beta_{\text{std}}=-0.073$), and the left superiorfrontal gyrus ($\beta_{\text{std}}=-0.066$). After Bonferroni correction for multiple testing, two regions – the left lateral orbitofrontal gyrus (corrected $p_{\text{SANS}}=0.034$) and left the pars opercularis (corrected $p_{\text{SANS}}=0.02$) - remained significant (SM section 2.2). The direction, size and lateralization of the effect compared well to our main results.

Discussion

Summary

In this study, we investigated the relationship between MOFC thinning and negative symptom severity in schizophrenia. We found that negative symptoms related inversely to cortical thickness in this brain region, with effects appearing greater in the left hemisphere compared to the right. This finding was independent of general illness severity, age and gender, but somewhat lessened when covarying for the influence of antipsychotic medication, age of onset and duration of illness (with the latter two variables being highly correlated). Exploratory analyses identified associations between negative symptoms and

thickness in the lateral orbitofrontal gyrus and pars opercularis in the left hemisphere. Our investigation has a number of strengths. First, using a meta-analytical approach, we were able to increase our sample size by a magnitude of 10 compared to previous studies. Second, the large sample allowed for the investigation of potentially influencing, but small effects of age, gender, illness severity, duration of illness, age of onset, antipsychotic medication, and handedness. Third, our sample included diverse patient groups that spans a broad age range allowing for a generalization of our findings.

Cortical thickness in the left medial orbitofrontal cortex and negative symptoms

Our results are in agreement with two previously published studies. Specifically, the association between MOFC thickness and negative symptoms in the study by Venkatasubramanian *et al.* (2008) was also negative and only present in the left hemisphere. Comparing patient groups with distinct symptom profiles, the study by Nenadic *et al.* (2015) also reported prefrontal reductions (including the MOFC) in patients characterized by predominantly negative symptoms compared to subgroups with predominantly paranoid and disorganised symptoms. However, there have also been reports, which failed to identify an association between MOFC thickness and negative symptoms (Xiao *et al.* 2013; Bodnar *et al.* 2014; Ansell *et al.* 2015; McKechnie *et al.* 2015). There are several potential explanations for these discrepant findings. First, all previous studies were based on rather small samples (between 40 and 130 patients) and hence

potentially underpowered to detect the weak, but robust association, we were able to identify in our study based on almost 2 000 patients. Furthermore, previous samples differed substantially in important characteristics such as illness chronicity and mean age. Of note, all studies that accepted the null hypothesis consisted of young (< 25 years), first-episode patients, while studies which reported effects, such as by Venkatasubramanian *et al.* (2008) and Nenadic *et al.* (2015), included older (>30 years), chronic patients. Similarly, mean age across our samples ranged from 28 – 43 years while mean duration of illness was 10 years, indicating that the association between negative symptoms and MOFC thickness might be more apparent when patients are older or in later stages of the disorder.

Animal and human studies suggest that neurons in the MOFC encode the subjective value of expected reward (Furuyashiki & Gallagher, 2007; Roesch & Olson, 2007), providing evidence that the MOFC mediates processes associated with the learning and retrieval of (subjective) value information, which guides decision-making and goal-directed behaviour (Bechara *et al.* 1994, 2000). Several studies suggest that negative symptoms in schizophrenia are associated with reinforcement learning abnormality (Waltz *et al.* 2007; Strauss *et al.* 2011). Gold *et al.* (2012) observed that patients with high-negative symptoms were less able to take expected reward values into account during decision making processes. As a result, patients fail to learn actions that lead to positive outcomes, while they outperform controls in avoiding punishing outcomes, showing that the findings were not indicative of general learning deficits, but of deficits in reward-related learning. Such

findings may help to explain the high prevalence of avolition and anhedonia in schizophrenia, two core dimensions of negative symptomatology. In line with this, aberrant neural responses to reward feedback was observed in medial prefrontal areas of patients with schizophrenia (Schlagenhauf *et al.* 2009), although results from this and several other studies also emphasize the relevance of ventral striatal pathways pointing towards an involvement of a wider striatal-prefrontal network in reward feedback (Schlagenhauf *et al.* 2014; Mørch-Johnsen *et al.* 2015; Radua *et al.* 2015).

The involvement of additional frontal brain regions is also supported by the results of our study, as exploratory analyses showed further associations of negative symptoms with the left lateral orbitofrontal gyrus and left pars opercularis. In line with our initial hypothesis, the lateral orbitofrontal gyrus is also predominantly involved in value processing (Kringelbach & Rolls, 2004; Zald *et al.* 2014). The pars opercularis is mainly linked to language processing (Belyk & Brown, 2014), which might relate to poverty of speech. An association of comparable effect size to our main finding, but not significant after Bonferroni correction, was also identified in the superior frontal gyrus (including the pre-supplementary motor area that is involved in volition (Haggard 2008; Bracht *et al.* 2013)), which might be associated with the lack of ability for spontaneous, self-generated action that relates strongly to the avolition subdomain of negative symptoms.

We observed a significant association between grey matter thickness and negative symptoms only in the left hemisphere. This aligns well with several other reports of

significant grey matter reductions in frontal regions of the left hemisphere (but not the right hemisphere) in patients with schizophrenia (Suzuki *et al.* 2002; Kawasaki *et al.* 2004; Honea *et al.* 2005; Kawasaki *et al.* 2008;). Moreover, cerebral lateralization of patients with schizophrenia tends to be less leftward lateralized and may even be rightward lateralized rather than symmetric. While some have argued that this might possibly reflect perturbations in the lateralization process underlying left cerebral dominance for language (Kawasaki *et al.* 2008), others believe that this atypical lateralization represents a greater involvement of the right hemisphere, which may relate to a broader, more diffuse semantic network (Grabner *et al.* 2007). With respect to value processing, the left prefrontal cortex has been linked stronger to motivation and positive affect than the right (Davidson, 2004; Price & Harmon-Jones, 2011). Hence, structural abnormalities in this region could link to deficits in the capacity to experience positive affect, a hallmark feature of negative symptoms.

In sum, our finding of a negative association between left MOFC thickness and negative symptoms underline the importance of this region in motivational and executive functioning, which is commonly impaired in schizophrenia patients (Barch & Dowd, 2010).

Potential moderators

Illness severity

We found that the effect of negative symptoms on left MOFC thickness was independent of general illness severity. That is, the MOFC might be specifically involved in motivational and executive aspects of schizophrenia as opposed to general schizophrenia psychopathology. Two other cortical thickness studies also failed to identify a significant correlation between illness severity (as measured using PANSS total scores) and thickness in any cortical region (Rimol *et al.* 2010; Oertel-Knöchel *et al.* 2013). Furthermore, van Haren *et al.* (2011) identified an association between *poor functional and symptomatic outcome* (derived using a factor analysis on Global Assessment of Functioning (GAF), the Camberwell Assessment of Need scale, and PANSS) and cortical thinning in the superior temporal gyrus, but not in frontal areas. Also in line with this is a functional imaging study by Simon *et al.* (2015), who investigated whole-brain activation during a reward anticipation paradigm. The authors observed that negative symptoms (especially signs of apathy), but not positive symptoms, were inversely correlated with activation in the ventral striatum, which at the same time showed reduced connectivity with the MOFC in patients compared to controls, further supporting the role of the MOFC in negative symptoms in schizophrenia.

Duration of illness

Duration of illness correlated strongly and negatively with left MOFC thickness (but not with negative symptom scores), although effects were dependent on age effects on thickness. Negative symptoms no longer predicted thickness significantly when covarying

for age *and* duration of illness (which was highly correlated with age and not predictive of MOFC thickness in the combined model) in the same model. This may indicate that some of the variance in thickness explained by negative symptoms depends on these variables. In this respect and as explained above, it is of interest that studies of first-episode patients did not find an association between symptoms and thickness in the MOFC. Ansell *et al.* (2015) studied very young (mean age of 22 years) psychosis patients, who had been diagnosed for the first time within the last three months. The authors found no effects of negative symptoms on thickness in prefrontal areas (although they did report that negative symptoms related to reduced thickness in parietal regions, but only in patients treated with first-generation antipsychotics). Crespo-Facorro *et al.* (2011) also studied first-episode patients and reported no significant relationship between negative symptoms and lobar cortical thickness. In the light of these previous findings and considering that participants in our study were predominantly older, chronic patients, our results might be more specific to later stages of the disorder.

Antipsychotic medication

Although the relationship between negative symptoms and left MOFC thickness did not vary by antipsychotic medication group in our moderator analysis, some amount of variance in left MOFC thickness was explained through antipsychotic medication effects, with patients treated with atypical antipsychotics showing the most robust reductions.

Notably, when we used current medication in CPZ equivalent units as a quantitative measures of antipsychotic medication (at the cost of a reduced sample size), an effect on MOFC thickness was not observed. While past studies predominantly reported volume and thickness deficits in frontal brain regions in patients treated with typical compared to those treated with atypical antipsychotics or unmedicated patients (van Haren *et al.* 2011; Lesh *et al.* 2015; Vita *et al.* 2015), there have also been many conflicting findings (Cahn *et al.* 2002; Ho *et al.* 2011). Attempting to reconcile previous reports, it has been suggested that antipsychotic medication effects on brain structure might be substance-specific (Xiao *et al.* 2008) and might also depend on illness chronicity and the duration of untreated psychosis. Some studies reported that medication effects on frontal brain structures were apparent especially during the early phase of treatment and that the effect size increased with longer durations of untreated psychosis (for a review see (Aderhold *et al.* 2014)). Genetic predisposition and substance use could also interact with antipsychotic medication (Tregellas *et al.* 2007; Hartz *et al.* 2010). With respect to our own findings we would like to stress that this is a purely correlational result, which does not in any way imply a causal relationship between antipsychotic medication intake and cortical thinning. Considering that we only investigated the possible effect of current antipsychotic medication, future studies are necessary to investigate this relationship in more detail by studying estimates of cumulative antipsychotic medication such as equivalent dose by time of exposure or treatment intensity in dose-years (Andreasen *et al.* 2010, 2013).

Limitations

The results of this study have to be seen in the light of the following limitations. First, our study was strongly hypothesis-driven in that we focussed on effects in the medial orbitofrontal cortex due to its role in value processing. As part of our exploratory analyses we also identified associations of comparable effect size in other brain regions including the pars opercularis and lateral orbitofrontal gyrus. Further studies are needed to investigate whether these associations are driven by processes that are not directly linked to value processing. Secondly, given the cross-sectional design of this study, we were unable to address directional effects. That is, we were not able to determine whether cortical thinning precedes or follows the development of negative symptoms. A range of factors such as measures of cumulative antipsychotic medication or relapse duration (Andreasen *et al.* 2010, 2013), which we were unable to investigate, might modulate the observed association and should be investigated in subsequent studies. Third, we included patients within the schizophrenia spectrum and used a measure of global negative symptom severity, but it is possible that effects were driven by schizophrenia subtype or subdimension-specific characteristics.

Conclusion

We investigated the relationship between negative symptoms and cortical thickness in the medial orbitofrontal cortex in a large cohort of schizophrenia patients, comprising almost 2 000 patients. Negative symptom severity was significantly associated with thickness in this region in the left, but not the right hemisphere. This association was irrespective of age, gender and illness severity, but possibly modulated by antipsychotic medication and duration of illness. Our findings provide further insight into symptom-related pathophysiological processes of schizophrenia.

Financial support

ENIGMA was supported in part by a Consortium grant (U54 EB020403 to PMT) from the NIH Institutes contributing to the Big Data to Knowledge (BD2K) Initiative, including the NIBIB and NCI. The authors would also like to express their gratitude for the personal support from the Deutsche Forschungsgemeinschaft (Research Fellowship to EW; Wa 3635/1-1). For additional support, see supplementary section 3.

Conflict of interest

The authors of this manuscript have no financial conflicts of interest to disclose.

Collaborators

Members of the Karolinska Schizophrenia Project consortium (KaSP): Farde L., Karolinska Institutet, Department of Clinical Neuroscience, Centre for Psychiatry Research, Stockholm, Sweden; Flyckt L., Karolinska Institutet, Department of Clinical Neuroscience, Centre for Psychiatry Research, Stockholm, Sweden; Engberg G., Karolinska Institutet, Department of Physiology and Pharmacology, Stockholm, Sweden; Erhardt S., Karolinska Institutet, Department of Physiology and Pharmacology, Stockholm, Sweden; Fatouros-Bergman H., Karolinska Institutet, Department of Clinical Neuroscience, Centre for Psychiatry Research, Stockholm, Sweden; Cervenka S., Karolinska Institutet, Department of Clinical Neuroscience, Centre for Psychiatry Research, Stockholm, Sweden; Schwieler L., Karolinska Institutet, Department of Physiology and Pharmacology, Stockholm, Sweden; Piehl F., Karolinska Institutet, Department of Clinical Neuroscience, Neuroimmunology Unit, Stockholm, Sweden; Agartz I., NORMENT, KG Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, University of Oslo, and Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway, and Department of Clinical Neuroscience, Karolinska Institutet, Centre for Psychiatry Research; Ikonen P., Karolinska Institutet, Department of Clinical Neuroscience, Centre for Psychiatry Research, Stockholm, Sweden; Collste K., Karolinska Institutet, Department of Clinical Neuroscience, Centre for Psychiatry Research, Stockholm, Sweden; Orhan F., Karolinska Institutet, Department of Physiology and Pharmacology, Stockholm, Sweden; Malmqvist A., Karolinska Institutet, Department of

Physiology and Pharmacology, Stockholm, Sweden; Hedberg M., Karolinska Institutet, Department of Physiology and Pharmacology, Stockholm, Sweden.

References

Aderhold DV, Weinmann S, Hägele C, Heinz A (2014). Frontale Hirnvolumenminderung durch Antipsychotika? *Der Nervenarzt* **86**, 302–323.

Andreasen NC (1983). Scale for the assessment of negative symptoms. *Iowa City, University of Iowa*

Andreasen NC, Liu D, Ziebell S, Vora A, Ho B-C (2013). Relapse duration, treatment intensity, and brain tissue loss in schizophrenia: a prospective longitudinal MRI study. *The American Journal of Psychiatry* **170**, 609–615.

Andreasen NC, Olsen S (1982). Negative v positive schizophrenia. Definition and validation. *Archives of General Psychiatry* **39**, 789–794.

Andreasen NC, Pressler M, Nopoulos P, Miller D, Ho B-C (2010). Antipsychotic dose equivalents and dose-years: a standardized method for comparing exposure to different drugs. *Biological Psychiatry* **67**, 255–262.

Ansell BRE, Dwyer DB, Wood SJ, Bora E, Brewer WJ, Proffitt TM, Velakoulis D, McGorry PD, Pantelis C (2015). Divergent effects of first-generation and second-generation

antipsychotics on cortical thickness in first-episode psychosis. *Psychological Medicine* **45**, 515–527.

Barch DM, Dowd EC (2010). Goal Representations and Motivational Drive in Schizophrenia: The Role of Prefrontal–Striatal Interactions. *Schizophrenia Bulletin* **36**, 919–934.

Bechara A, Damasio AR, Damasio H, Anderson SW (1994). Insensitivity to future consequences following damage to human prefrontal cortex. *Cognition* **50**, 7–15.

Bechara A, Tranel D, Damasio H (2000). Characterization of the decision-making deficit of patients with ventromedial prefrontal cortex lesions. *Brain* **123**, 2189–2202.

Behrendt S (2014). *lm.beta: Add Standardized Regression Coefficients to lm-Objects*

Belyk M, Brown S (2014). Perception of affective and linguistic prosody: an ALE meta-analysis of neuroimaging studies. *Social Cognitive and Affective Neuroscience* **9**, 1395–1403.

Berridge KC, Kringelbach ML (2015). Pleasure systems in the brain. *Neuron* **86**, 646–664.

Bodnar M, Hovington CL, Buchy L, Malla AK, Joober R, Lepage M (2014). Cortical Thinning in Temporo-Parietal Junction (TPJ) in Non-Affective First-Episode of Psychosis Patients with Persistent Negative Symptoms. *PLoS ONE* **9**, e101372.

Bracht T, Schnell S, Federspiel A, Razavi N, Horn H, Strik W, Wiest R, Dierks T, Müller TJ, Walther S (2013). Altered cortico-basal ganglia motor pathways reflect reduced volitional motor activity in schizophrenia. *Schizophrenia Research* **143**, 269–276.

Burgdorf J, Panksepp J (2006). The neurobiology of positive emotions. *Neuroscience & Biobehavioral Reviews* **30**, 173–187.

Cahn W, Hulshoff Pol HE, Lems EBTE, van Haren NEM, Schnack HG, van der Linden JA, Schothorst PF, van Engeland H, Kahn RS (2002). Brain volume changes in first-episode schizophrenia: a 1-year follow-up study. *Archives of General Psychiatry* **59**, 1002–1010.

Cohen AS, Minor KS (2010). Emotional Experience in Patients With Schizophrenia Revisited: Meta-analysis of Laboratory Studies. *Schizophrenia Bulletin* **36**, 143–150.

Crespo-Facorro B, Roiz-Santiañez R, Pérez-Iglesias R, Rodriguez-Sanchez JM, Mata I, Tordesillas-Gutierrez D, Sanchez E, Tabarés-Seisdedos R, Andreasen N, Magnotta V, Vázquez-Barquero JL (2011). Global and regional cortical thinning in first-episode psychosis patients: relationships with clinical and cognitive features. *Psychological Medicine* **41**, 1449–1460.

Davidson RJ (2004). Well-being and affective style: neural substrates and biobehavioural correlates. *Philosophical Transactions-Royal Society of London Series B Biological Sciences*, 1395–1412.

Deserno L, Boehme R, Heinz A, Schlagenhauf F (2013). Reinforcement Learning and Dopamine in Schizophrenia: Dimensions of Symptoms or Specific Features of a Disease Group? *Frontiers in Psychiatry* **4**, 172.

Deserno L, Schlagenhauf F, Heinz A (2016). Striatal dopamine, reward, and decision making in schizophrenia. *Dialogues in Clinical Neuroscience* **18**, 77–89.

Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM, Maguire RP, Hyman BT, Albert MS, Killiany RJ (2006). An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage* **31**, 968–980.

Ducharme S, Albaugh MD, Hudziak JJ, Botteron KN, Nguyen T-V, Truong C, Evans AC, Karama S, Ball WS, Byars AW, Schapiro M, Bommer W, Carr A, German A, Dunn S, Rivkin MJ, Waber D, Mulkern R, Vajapeyam S, Chiverton A, Davis P, Koo J, Marmor J, Mrakotsky C, Robertson R, McAnulty G, Brandt ME, Fletcher JM, Kramer LA, Yang G, McCormack C, Hebert KM, Volero H, Botteron K, McKinstry RC, Warren W, Nishino T, Almli CR, Todd R, Constantino J, McCracken JT, Levitt J, Alger J, O'Neil J, Toga A, Asarnow R, Fadale D, Heinichen L, Ireland C, Wang D-J, Moss E, Zimmerman RA, Bintliff B, Bradford R, Newman J, Evans AC, Arnaoutelis R, Pike GB, Collins DL, Leonard G, Paus T, Zijdenbos A, Das S, Fonov V, Fu L, Harlap J, Leppert I, Milovan D, Vins D, Zeffiro T, Meter JV, Lange N, Froimowitz MP, Botteron K, Almli CR, Rainey C, Henderson S, Nishino T, Warren W, Edwards JL, Dubois D, Smith K, Singer T, Wilber AA, Pierpaoli C, Bassar PJ, Chang L-C, Koay CG, Walker L, Freund L, Rumsey J, Baskir L, Stanford L, Sirocco K, Gwinn-Hardy K, Spinella G, McCracken JT, Alger JR, Levitt J, O'Neill J. (2014). Anxious/Depressed Symptoms are

Linked to Right Ventromedial Prefrontal Cortical Thickness Maturation in Healthy Children and Young Adults. *Cerebral Cortex* **24**, 2941–2950.

Ehrlich S, Brauns S, Yendiki A, Ho B-C, Calhoun V, Schulz SC, Gollub RL, Sponheim SR (2012). Associations of cortical thickness and cognition in patients with schizophrenia and healthy controls. *Schizophrenia bulletin* **38**, 1050–1062.

van Erp TGM, Hibar DP, Rasmussen JM, Glahn DC, Pearlson GD, Andreassen OA, Agartz I, Westlye LT, Haukvik UK, Dale AM, Melle I, Hartberg CB, Gruber O, Kraemer B, Zilles D, Donohoe G, Kelly S, McDonald C, Morris DW, Cannon DM, Corvin A, Machielsen MWJ, Koenders L, de Haan L, Veltman DJ, Satterthwaite TD, Wolf DH, Gur RC, Gur RE, Potkin SG, Mathalon DH, Mueller BA, Preda A, Macciardi F, Ehrlich S, Walton E, Hass J, Calhoun VD, Bockholt HJ, Sponheim SR, Shoemaker JM, van Haren NEM, Pol HEH, Ophoff RA, Kahn RS, Roiz-Santiañez R, Crespo-Facorro B, Wang L, Alpert KI, Jönsson EG, Dimitrova R, Bois C, Whalley HC, McIntosh AM, Lawrie SM, Hashimoto R, Thompson PM, Turner JA (2015). Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Molecular Psychiatry* **21**, 547-553.

van Erp TGM, Preda A, Nguyen D, Faziola L, Turner J, Bustillo J, Belger A, Lim KO, McEwen S, Voyvodic J, Mathalon DH, Ford J, Potkin SG, Fbirn null (2014). Converting positive and negative symptom scores between PANSS and SAPS/SANS. *Schizophrenia Research* **152**, 289–294.

Fellows LK, Farah MJ (2005). Different Underlying Impairments in Decision-making Following Ventromedial and Dorsolateral Frontal Lobe Damage in Humans. *Cerebral Cortex* **15**, 58–63.

Fisher T, Shamay-Tsoory SG, Eran A, Aharon-Peretz J (2011). Characterization of recovery and neuropsychological consequences of orbitofrontal lesion: A case study. *Neurocase* **17**, 285–293.

Furuyashiki T, Gallagher M (2007). Neural encoding in the orbitofrontal cortex related to goal-directed behavior. *Annals of the New York Academy of Sciences* **1121**, 193–215.

Geisler D, Walton E, Naylor M, Roessner V, Lim KO, Charles Schulz S, Gollub RL, Calhoun VD, Sponheim SR, Ehrlich S (2015). Brain structure and function correlates of cognitive subtypes in schizophrenia. *Psychiatry Research* **234**, 74–83.

Gold JM, Waltz JA, Matveeva TM, Kasanova Z, Strauss GP, Herbener ES, Collins AGE, Frank MJ (2012). Negative symptoms and the failure to represent the expected reward value of actions: Behavioral and computational modeling evidence. *Archives of General Psychiatry* **69**, 129–138.

Goldman AL (2009). Widespread Reductions of Cortical Thickness in Schizophrenia and Spectrum Disorders and Evidence of Heritability. *Archives of General Psychiatry* **66**, 467.

Grabenhorst F, Rolls ET (2011). Value, pleasure and choice in the ventral prefrontal cortex.

Trends in Cognitive Sciences **15**, 56–67.

Grabner RH, Fink A, Neubauer AC (2007). Brain correlates of self-rated originality of ideas: evidence from event-related power and phase-locking changes in the EEG. *Behavioral Neuroscience* **121**, 224–230.

Haggard P (2008). Human volition: towards a neuroscience of will. *Nature Reviews Neuroscience* **9**, 934–946.

Hartz SM, Ho B-C, Andreasen NC, Librant A, Rudd D, Epping EA, Wassink TH (2010). G72 influences longitudinal change in frontal lobe volume in schizophrenia. *American Journal of Medical Genetics. Part B, Neuropsychiatric Genetics: The Official Publication of the International Society of Psychiatric Genetics* **153B**, 640–647.

Harvey P-O, Armony J, Malla A, Lepage M (2010). Functional neural substrates of self-reported physical anhedonia in non-clinical individuals and in patients with schizophrenia. *Journal of Psychiatric Research* **44**, 707–716.

Harvey P-O, Pruessner J, Czechowska Y, Lepage M (2007). Individual differences in trait anhedonia: a structural and functional magnetic resonance imaging study in non-clinical subjects. *Molecular Psychiatry* **12**, 767–775.

Ho B, Andreasen N, Ziebell S, Pierson R, Magnotta V (2011). Long-term antipsychotic treatment and brain volumes: A longitudinal study of first-episode schizophrenia. *Archives of General Psychiatry* **68**, 128–137.

Honea R, Crow TJ, Passingham D, Mackay CE (2005). Regional Deficits in Brain Volume in Schizophrenia: A Meta-Analysis of Voxel-Based Morphometry Studies. *Am J Psychiatry* **162**, 2233–2245.

Hornak J, Bramham J, Rolls ET, Morris RG, O'Doherty J, Bullock PR, Polkey CE (2003). Changes in emotion after circumscribed surgical lesions of the orbitofrontal and cingulate cortices. *Brain: A Journal of Neurology* **126**, 1691–1712.

Jablensky A (2006). Subtyping schizophrenia: implications for genetic research. *Molecular Psychiatry* **11**, 815–836.

Jeppesen P, Petersen L, Thorup A, Abel M-B, Øhlenschläger J, Christensen TØ, Krarup G, Jørgensen P, Nordentoft M (2008). The association between pre-morbid adjustment, duration of untreated psychosis and outcome in first-episode psychosis. *Psychological Medicine* **38**, 1157–1166.

Kalkstein S, Hurford I, Gur RC (2010). Neurocognition in schizophrenia. *Current Topics in Behavioral Neurosciences* **4**, 373–390.

Kawasaki Y, Suzuki M, Nohara S, Hagino H, Takahashi T, Matsui M, Yamashita I, Chitnis XA, McGuire PK, Seto H, Kurachi M (2004). Structural brain differences in patients with schizophrenia and schizotypal disorder demonstrated by voxel-based morphometry. *European Archives of Psychiatry and Clinical Neuroscience* **254**, 406–414.

Kawasaki Y, Suzuki M, Takahashi T, Nohara S, McGuire PK, Seto H, Kurachi M (2008). Anomalous Cerebral Asymmetry in Patients with Schizophrenia Demonstrated by Voxel-Based Morphometry. *Biological Psychiatry* **63**, 793–800.

Kay SR, Fiszbein A, Opler LA (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia bulletin* **13**, 261–276.

Kringelbach ML, Rolls ET (2004). The functional neuroanatomy of the human orbitofrontal cortex: evidence from neuroimaging and neuropsychology. *Progress in Neurobiology* **72**, 341–372.

Kühn S, Gallinat J (2012). The neural correlates of subjective pleasantness. *NeuroImage* **61**, 289–294.

Lesh TA, Tanase C, Geib BR, Niendam TA, Yoon JH, Minzenberg MJ, Ragland JD, Solomon M, Carter CS (2015). A multimodal analysis of antipsychotic effects on brain structure and function in first-episode schizophrenia. *JAMA psychiatry* **72**, 226–234.

Liu X, Hairston J, Schrier M, Fan J (2011). Common and distinct networks underlying reward valence and processing stages: A meta-analysis of functional neuroimaging studies.

Neuroscience & Biobehavioral Reviews **35**, 1219–1236.

McKechanie AG, Moorhead TWJ, Stanfield AC, Whalley HC, Johnstone EC, Lawrie SM,

Owens DGC (2015). Negative symptoms and longitudinal grey matter tissue loss in adolescents at risk of psychosis: preliminary findings from a 6-year follow-up study. *The British Journal of Psychiatry* **6**, 565-70.

Milev P, Ho B-C, Arndt S, Andreasen NC (2005). Predictive Values of Neurocognition and Negative Symptoms on Functional Outcome in Schizophrenia: A Longitudinal First-Episode Study With 7-Year Follow-Up. *American Journal of Psychiatry* **162**, 495–506.

Mørch-Johnsen L, Nesvåg R, Faerden A, Haukvik UK, Jørgensen KN, Lange EH, Andreassen OA, Melle I, Agartz I (2015). Brain structure abnormalities in first-episode psychosis patients with persistent apathy. *Schizophrenia Research* **164**, 59–64.

Nenadic I, Yotter RA, Sauer H, Gaser C (2015). Patterns of cortical thinning in different subgroups of schizophrenia. *The British Journal of Psychiatry: The Journal of Mental Science* **206**, 479–483.

Nesvåg R, Lawyer G, Varnäs K, Fjell AM, Walhovd KB, Frigessi A, Jönsson EG, Agartz I (2008). Regional thinning of the cerebral cortex in schizophrenia: effects of diagnosis, age and antipsychotic medication. *Schizophrenia research* **98**, 16–28.

Oertel-Knöchel V, Knöchel C, Rotarska-Jagiela A, Reinke B, Prvulovic D, Haenschel C,

Hampel H, Linden DEJ (2013). Association between Psychotic Symptoms and Cortical

Thickness Reduction across the Schizophrenia Spectrum. *Cerebral Cortex* **23**, 61–70.

Pessoa L (2009). How do emotion and motivation direct executive control? *Trends in*

Cognitive Sciences **13**, 160–166.

Peters J, Büchel C (2010). Neural representations of subjective reward value. *Behavioural*

Brain Research **213**, 135–141.

Plailly J, d'Amato T, Saoud M, Royet J-P (2006). Left temporo-limbic and orbital dysfunction

in schizophrenia during odor familiarity and hedonicity judgments. *NeuroImage* **29**, 302–

313.

Price TF, Harmon-Jones E (2011). Approach motivational body postures lean toward left

frontal brain activity. *Psychophysiology* **48**, 718–722.

R Development Core Team (2008). *R: A language and environment for statistical computing*.

R Foundation for Statistical Computing. Vienna, Austria.

Radua J, Schmidt A, Borgwardt S, Heinz A, Schlagenhauf F, McGuire P, Fusar-Poli P (2015).

Ventral Striatal Activation During Reward Processing in Psychosis: A Neurofunctional Meta-

Analysis. *JAMA psychiatry* **72**, 1243–1251.

- Rimol LM, Hartberg CB, Nesvåg R, Fennema-Notestine C, Hagler Jr. DJ, Pung CJ, Jennings RG, Haukvik UK, Lange E, Nakstad PH, Melle I, Andreassen OA, Dale AM, Agartz I** (2010). Cortical Thickness and Subcortical Volumes in Schizophrenia and Bipolar Disorder. *Biological Psychiatry* **68**, 41–50.
- Roesch MR, Olson CR** (2007). Neuronal activity related to anticipated reward in frontal cortex: does it represent value or reflect motivation? *Annals of the New York Academy of Sciences* **1121**, 431–446.
- Rosenheck R, Leslie D, Keefe R, McEvoy J, Swartz M, Perkins D, Stroup S, Hsiao JK, Lieberman J** (2006). Barriers to Employment for People With Schizophrenia. *American Journal of Psychiatry* **163**, 411–417.
- Schlagenhauf F, Huys QJM, Deserno L, Rapp MA, Beck A, Heinze H-J, Dolan R, Heinz A** (2014). Striatal dysfunction during reversal learning in unmedicated schizophrenia patients. *NeuroImage* **89**, 171–180.
- Schlagenhauf F, Sterzer P, Schmack K, Ballmaier M, Rapp M, Wrase J, Juckel G, Gallinat J, Heinz A** (2009). Reward Feedback Alterations in Unmedicated Schizophrenia Patients: Relevance for Delusions. *Biological Psychiatry* **65**, 1032–1039.
- Schultz CC, Koch K, Wagner G, Roebel M, Schachtzabel C, Gaser C, Nenadic I, Reichenbach JR, Sauer H, Schlösser RGM** (2010). Reduced cortical thickness in first episode schizophrenia. *Schizophrenia Research* **116**, 204–209.

Segarra N, Metastasio A, Ziauddeen H, Spencer J, Reinders NR, Dudas RB, Arrondo G, Robbins TW, Clark L, Fletcher PC, Murray GK (2016). Abnormal Frontostriatal Activity During Unexpected Reward Receipt in Depression and Schizophrenia: Relationship to Anhedonia. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology* **41**, 2001–2010.

Simon JJ, Cordeiro SA, Weber M-A, Friederich H-C, Wolf RC, Weisbrod M, Kaiser S (2015). Reward System Dysfunction as a Neural Substrate of Symptom Expression Across the General Population and Patients With Schizophrenia. *Schizophrenia Bulletin* **41**, 1370-1378.

Strauss GP, Frank MJ, Waltz JA, Kasanova Z, Herbener ES, Gold JM (2011). Deficits in positive reinforcement learning and uncertainty-driven exploration are associated with distinct aspects of negative symptoms in schizophrenia. *Biological Psychiatry* **69**, 424–431.

Suzuki M, Nohara S, Hagino H, Kurokawa K, Yotsutsuji T, Kawasaki Y, Takahashi T, Matsui M, Watanabe N, Seto H, Kurachi M (2002). Regional changes in brain gray and white matter in patients with schizophrenia demonstrated with voxel-based analysis of MRI. *Schizophrenia Research* **55**, 41–54.

Szatkowska I, Bogorodzki P, Wolak T, Marchewka A, Szeszkowski W (2008). The effect of motivation on working memory: An fMRI and SEM study. *Neurobiology of Learning and Memory* **90**, 475–478.

Takeuchi H, Taki Y, Nouchi R, Sekiguchi A, Kotozaki Y, Miyauchi CM, Yokoyama R, Iizuka K, Hashizume H, Nakagawa S, Kunitoki K, Sassa Y, Kawashima R (2014). Regional gray matter density is associated with achievement motivation: evidence from voxel-based morphometry. *Brain Structure & Function* **219**, 71–83.

Tregellas JR, Shatti S, Tanabe JL, Martin LF, Gibson L, Wylie K, Rojas DC (2007). Gray matter volume differences and the effects of smoking on gray matter in schizophrenia. *Schizophrenia Research* **97**, 242–249.

van Haren NM, Schnack HG, Cahn W, van den Heuvel MP, Lepage C, Collins L, Evans AC, Hulshoff Pol HE, Kahn RS (2011). Changes in cortical thickness during the course of illness in schizophrenia. *Archives of General Psychiatry* **68**, 871–880.

Venkatasubramanian G, Jayakumar PN, Gangadhar BN, Keshavan MS (2008). Automated MRI parcellation study of regional volume and thickness of prefrontal cortex (PFC) in antipsychotic-naïve schizophrenia. *Acta Psychiatrica Scandinavica* **117**, 420–431.

Viechtbauer (2010). Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software* **36**, 1–48.

Vita A, De Peri L, Deste G, Barlati S, Sacchetti E (2015). The Effect of Antipsychotic Treatment on Cortical Gray Matter Changes in Schizophrenia: Does the Class Matter? A Meta-analysis and Meta-regression of Longitudinal Magnetic Resonance Imaging Studies. *Biological Psychiatry* **78**, 403–412.

Waltz JA, Frank MJ, Robinson BM, Gold JM (2007). Selective reinforcement learning deficits in schizophrenia support predictions from computational models of striatal-cortical dysfunction. *Biological Psychiatry* **62**, 756–764.

Woods SW (2003). Chlorpromazine equivalent doses for the newer atypical antipsychotics. *The Journal of Clinical Psychiatry* **64**, 663–667.

Xiao L, Xu H, Zhang Y, Wei Z, He J, Jiang W, Li X, Dyck LE, Devon RM, Deng Y, Li XM (2008). Quetiapine facilitates oligodendrocyte development and prevents mice from myelin breakdown and behavioral changes. *Molecular Psychiatry* **13**, 697–708.

Xiao Y, Lui S, Deng W, Yao L, Zhang W, Li S, Wu M, Xie T, He Y, Huang X, Hu J, Bi F, Li T, Gong Q (2015). Altered Cortical Thickness Related to Clinical Severity But Not the Untreated Disease Duration in Schizophrenia. *Schizophrenia Bulletin* **41**, 201–210.

Zald DH, McHugo M, Ray KL, Glahn DC, Eickhoff SB, Laird AR (2014). Meta-Analytic Connectivity Modeling Reveals Differential Functional Connectivity of the Medial and Lateral Orbitofrontal Cortex. *Cerebral Cortex* **24**, 232–248.

Table

	estimate	range	data available for N number of studies
% males	68	55-76	17
Mean age in years	34	28-43	17
Mean SANS Global	7.91	2.86-12.90	17
Mean duration of illness in years	10	1-20	13
Mean Age of Onset in years	24	19-29	13
Mean illness severity (PANSS Total)	70.43	49.81-90.22	9
Antipsychotic medication			13
%Atypical (N)	71 (1201)	39-91	
%Typical (N)	11 (202)	0-45	
%Both A & T (N)	9 (157)	0-24	
%None (N)	9 (149)	0-53	
Mean Chlorpromazine equivalents	423.32	97.48-637.80	12
Handedness			14
%Right	90	68-95	
%Left	8	2-14	
%Ambidextrous	2	0-25	
Cortical thickness in mm			17
Mean left medial orbitofrontal	2.46	2.26-2.71	
Mean right medial orbitofrontal	2.42	2.17-2.61	

Table 1. Demographics. Means are weighted by study sample size.

Figure legend

Figure 1. Forest plot of association between global SANS and cortical thickness in the left medial orbitofrontal cortex across all 17 study sites, controlling for age, gender and number of sites (if applicable; of note the global SANS-left MOFC thickness relationship did not differ between single vs multisite samples ($p=0.422$; see supplementary section 2.1, subsection o)). Fisher's transformed standardized regression coefficients are denoted by black boxes. Black lines indicate 95% confidence intervals. The combined estimate for all sites is represented by a black diamond with the outer edges of the diamond indicating the confidence interval limits.

Supplementary Materials


SM 1 Method*SM 1.1 Symptom score conversion algorithms*

Global SANS (Summary) Score = $-2.0671 + (0.665 * \text{PANSS Negative score})$

Global SANS (Summary) Score = $1.0863 + (0.2943 * \text{Total SANS (composite) score})$

SM 1.2 Imaging descriptives and quality control

Parcellations were visually inspected and statistically evaluated for outliers following standardized ENIGMA protocols (<http://enigma.ini.usc.edu/protocols/imaging-protocols>). Each image parcellation was individually examined by a neuroimaging expert at each site by overlaying the parcellation label of each structure on the T1-weighted brain scan. Further, we collected study-wide statistics (means, minimums, maximums, and standard deviations; see SM Figure 1s) as well as histogram and diagnostic plots in order to identify non-normally distributed data and major outliers. A subject was considered a statistical outlier if its thickness was >2.698 standard deviations away from the global mean. For each subject that was marked as a statistical outlier, individual sites were asked to re-inspect the subject's parcellations in order to verify that it was properly segmented. If a subject was a statistical outlier, but was properly segmented it was kept in the analysis. Otherwise the subject was removed (see SM Table 2s for details).



SM Figure 1s. Left (A-B) and right (C-D) medial orbitofrontal cortical thickness descriptives (min/mean/sd/max) by study site.

SM 2 Results

SM 2.1 Detailed results of meta-analyses

a) Main model: left MOFC <- global SANS + gender + age + site (if applicable)

In the main model we investigated the effect of global SANS scores on left MOFC thickness, covarying for gender, age and the number of sites as dummy variables (where applicable). Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of the global SANS score on left MOFC thickness. Results were based on estimates from 17 studies using a restricted maximum likelihood (REML) approach. Also provided are estimates and significance levels for effect size heterogeneity between studies.

Random-Effects Model (k = 17; tau² estimator: REML)

tau² (estimated amount of total heterogeneity): 0.0067 (SE = 0.0057)

tau (square root of estimated tau² value): 0.0816

I² (total heterogeneity / total variability): 42.38%


H² (total variability / sampling variability): 1.74

Test for Heterogeneity:

Q(df = 16) = 26.3994, p-val = 0.0487

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub	
-0.0752	0.0321	-2.3446	0.0190	-0.1381	-0.0123	*

Funnel plot main model

SM Figure 2s. Funnel plot. Individual study regression coefficients are plotted against sample variance (a measure of the precision of the data).

b) Exploratory model: right MOFC <- global SANS + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of the global SANS score on right MOFC thickness. Results were based on estimates from 17 studies.

Random-Effects Model (k = 17; tau² estimator: REML)

tau² (estimated amount of total heterogeneity): 0.0080 (SE = 0.0063)

tau (square root of estimated tau² value): 0.0892

I² (total heterogeneity / total variability): 46.78%

H² (total variability / sampling variability): 1.88

Test for Heterogeneity:

Q(df = 16) = 28.4025, p-val = 0.0283

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub	
-0.0643	0.0335	-1.9207	0.0548	-0.1299	0.0013	.

c) Effects of covariates: left MOFC <- global SANS + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of age on left MOFC thickness. Results were based on estimates from 17 studies.

Random-Effects Model (k = 17; tau² estimator: REML)

tau² (estimated amount of total heterogeneity): 0.0117 (SE = 0.0078)

tau (square root of estimated tau² value): 0.1083

I² (total heterogeneity / total variability): 56.43%

H² (total variability / sampling variability): 2.30

Test for Heterogeneity:

$Q(df = 16) = 36.1039$, $p\text{-val} = 0.0028$

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub	
-0.2372	0.0371	-6.3925	<.0001	-0.3100	-0.1645	***

d) Effects of covariates: left MOFC <- global SANS + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of gender on left MOFC thickness.

Results were based on estimates from 17 studies.

Random-Effects Model (k = 17; tau² estimator: REML)

tau² (estimated amount of total heterogeneity): 0 (SE = 0.0026)

tau (square root of estimated tau² value): 0

I² (total heterogeneity / total variability): 0.00%

H² (total variability / sampling variability): 1.00

Test for Heterogeneity:

$Q(df = 16) = 13.5923$, $p\text{-val} = 0.6291$

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub	
-0.0900	0.0227	-3.9592	<.0001	-0.1346	-0.0455	***

- e) Effects of covariates: left MOFC <- global SANS + illness severity + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of illness severity on left MOFC thickness. Results were based on estimates from 9 studies.

Random-Effects Model (k = 9; tau² estimator: REML)

tau² (estimated amount of total heterogeneity): 0.0387 (SE = 0.0254)

tau (square root of estimated tau² value): 0.1967

I² (total heterogeneity / total variability): 81.26%

H² (total variability / sampling variability): 5.34

Test for Heterogeneity:

Q(df = 8) = 44.9511, p-val < .0001

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub
0.0446	0.0755	0.5908	0.5546	-0.1034	0.1927

- f) Effects of covariates: left MOFC <- global SANS + illness severity + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of global SANS score on left MOFC thickness. Results were based on estimates from 9 studies.

Random-Effects Model (k = 9; tau² estimator: REML)

tau² (estimated amount of total heterogeneity): 0.0145 (SE = 0.0126)

tau (square root of estimated tau² value): 0.1205

I² (total heterogeneity / total variability): 61.91%

H² (total variability / sampling variability): 2.63

Test for Heterogeneity:

Q(df = 8) = 21.8200, p-val = 0.0053

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub	
-0.1126	0.0536	-2.0998	0.0357	-0.2177	-0.0075	*

- g) Effects of covariates: left MOFC <- global SANS + length of illness + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of length of illness estimates on left MOFC thickness. Results were based on estimates from 13 studies.

Random-Effects Model (k = 13; tau² estimator: REML)

tau² (estimated amount of total heterogeneity): 0.0539 (SE = 0.0271)

tau (square root of estimated tau² value): 0.2321

I² (total heterogeneity / total variability): 86.66%

H² (total variability / sampling variability): 7.50

Test for Heterogeneity:

$Q(df = 12) = 51.8783, p\text{-val} < .0001$

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub
-0.0922	0.0719	-1.2836	0.1993	-0.2331	0.0486

- h) Effects of covariates: left MOFC <- global SANS + length of illness + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of global SANS scores on left MOFC thickness. Results were based on estimates from 13 studies.

Random-Effects Model (k = 13; tau² estimator: REML)

tau² (estimated amount of total heterogeneity): 0.0079 (SE = 0.0069)

tau (square root of estimated tau² value): 0.0892

I² (total heterogeneity / total variability): 48.94%

H² (total variability / sampling variability): 1.96

Test for Heterogeneity:

$Q(df = 12) = 22.8334, p\text{-val} = 0.0292$

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub
-0.0681	0.0373	-1.8240	0.0682	-0.1413	0.0051

- i) Moderation analysis of age of onset on the association between global SANS scores and left MOFC thickness

Shown below are the estimate, the standard error, p value and the confidence intervals for the moderating effects of age of onset on the meta-analytical standardized regression beta of global SANS scores on left MOFC thickness. Results were based on estimates from 13 studies.

Mixed-Effects Model ($k = 13$; τ^2 estimator: REML)

τ^2 (estimated amount of residual heterogeneity): 0.0077 (SE = 0.0072)

τ (square root of estimated τ^2 value): 0.0875

I^2 (residual heterogeneity / unaccounted variability): 46.82%

H^2 (unaccounted variability / sampling variability): 1.88

R^2 (amount of heterogeneity accounted for): 0.00%

Test for Residual Heterogeneity:

$QE(df = 11) = 20.0562$, $p\text{-val} = 0.0446$

Test of Moderators (coefficient(s) 2):

$QM(df = 1) = 0.1133$, $p\text{-val} = 0.7365$

Model Results:

	se	zval	pval	ci.lb	ci.ub
intrcpt	-0.0651	0.0370	-1.7578	0.0788	-0.1377 0.0075 .
l(AO_mean - 23.83)	0.0050	0.0149	0.3365	0.7365	-0.0242 0.0342

j) Moderation analysis of antipsychotic medication on the association between global SANS scores and left MOFC thickness

To investigate potentially moderating effects of antipsychotic medication, we derived the percentages of patients treated with typical or atypical antipsychotics, both antipsychotic medication types and the percentage of unmedicated patients in each sample. Shown below are the estimate, the standard error, p value and the confidence intervals for the moderating effects of antipsychotic medication on the meta-analytical standardized regression beta of global SANS scores on left MOFC thickness. Percentages of participants on typical, atypical and combined antipsychotic medication were mean centered and compared to the percentage of unmedicated patients (baseline). The intercept refers to the effect of global SANS score on left MOFC thickness after accounting for antipsychotic medication. Results were based on estimates from 13 studies.

Mixed-Effects Model (k = 13; tau² estimator: REML)

tau² (estimated amount of residual heterogeneity): 0.0039 (SE = 0.0056)
tau (square root of estimated tau² value): 0.0621
I² (residual heterogeneity / unaccounted variability): 32.04%
H² (unaccounted variability / sampling variability): 1.47

to the effect of global SANS score on left MOFC thickness after accounting for handedness. Results were based on estimates from 14 studies.

Mixed-Effects Model (k = 14; tau^2 estimator: REML)

tau^2 (estimated amount of residual heterogeneity):	0.0061 (SE = 0.0059)
tau (square root of estimated tau^2 value):	0.0780
I^2 (residual heterogeneity / unaccounted variability):	44.65%
H^2 (unaccounted variability / sampling variability):	1.81
R^2 (amount of heterogeneity accounted for):	0.00%

Test for Residual Heterogeneity:

QE(df = 11) = 19.0779, p-val = 0.0597

Test of Moderators (coefficient(s) 2,3):

QM(df = 2) = 1.0884, p-val = 0.5803

Model Results:

	estimate	se	zval	pval	ci.lb	ci.ub
intrcpt	-0.0603	0.0342	-1.7634	0.0778	-0.1273	0.0067
.						
I(HAND_a_perc - 3.591)	0.0010	0.0074	0.1399	0.8888	-0.0134	0.0154
I(HAND_l_perc - 8.688)	-0.0113	0.0116	-0.9744	0.3299	-0.0341	0.0115

N.B.: predictor abbreviations refer to the percentage of ambidextrous (HAND_a_perc), left-handed (HAND_l_perc) and right-handed patients (intrcpt, i.e. baseline).

l) Left MOFC thickness differences by antipsychotic medication type percentages

We also investigated whether left MOFC thickness differed by the percentage of patients treated with typical or atypical antipsychotics, both antipsychotic medication types and the percentage of unmedicated patients. Shown below are the estimate, the standard error, p value and the confidence intervals of mean left MOFC thickness for each antipsychotic medication group. Percentages were mean centered and compared to unmedicated patients (baseline). Results were based on estimates from 13 studies.

Mixed-Effects Model (k = 13; tau^2 estimator: REML)

tau^2 (estimated amount of residual heterogeneity):	0.0060 (SE = 0.0030)
tau (square root of estimated tau^2 value):	0.0777
I^2 (residual heterogeneity / unaccounted variability):	95.96%
H^2 (unaccounted variability / sampling variability):	24.73
R^2 (amount of heterogeneity accounted for):	44.87%

Test for Residual Heterogeneity:

QE(df = 9) = 249.4885, p-val < .0001

Test of Moderators (coefficient(s) 2,3,4):

QM(df = 3) = 12.2740, p-val = 0.0065

Model Results:

	estimate	se	zval	pval	ci.lb	ci.ub
intrcpt	2.4841	0.0222	111.8192	<.0001	2.4405	2.5276

I(MED_a_perc - 68.72)	-0.0058	0.0019	-3.1327	0.0017	-0.0094	-0.0022
**						
I(MED_b_perc - 8.536)	-0.0026	0.0033	-0.7653	0.4441	-0.0091	0.0040
I(MED_t_perc - 11.1)	-0.0065	0.0022	-2.9253	0.0034	-0.0109	-0.0021
**						

N.B.: predictor abbreviations refer to percentage of patients treated with atypical (MED_a_perc) or typical (MED_t_perc) antipsychotics, both antipsychotic medication types (MED_b_perc) and the percentage of unmedicated patients (intrcpt, i.e. baseline).

m) Global SANS score differences by antipsychotic medication type percentages

Furthermore, We also investigated whether global SANS scores differed by the percentage of patients treated with typical or atypical antipsychotics, both antipsychotic medication types and the percentage of unmedicated patients. Shown below are the estimate, the standard error, p value and the confidence intervals of mean global SANS scores for each antipsychotic medication group. Percentages were mean centered and compared to unmedicated patients (baseline). Results were based on estimates from 13 studies.

```
Mixed-Effects Model (k = 13; tau^2 estimator: REML)

tau^2 (estimated amount of residual heterogeneity):      7.3158 (SE = 3.5444)
tau (square root of estimated tau^2 value):             2.7048
I^2 (residual heterogeneity / unaccounted variability): 98.34%
H^2 (unaccounted variability / sampling variability):    60.25
R^2 (amount of heterogeneity accounted for):             0.00%

Test for Residual Heterogeneity:
QE(df = 9) = 483.0582, p-val < .0001

Test of Moderators (coefficient(s) 2,3,4):
QM(df = 3) = 2.6231, p-val = 0.4535

Model Results:
```


	estimate	se	zval	pval	ci.lb	ci.ub
intrcpt ***	8.7936	0.7616	11.5460	<.0001	7.3009	10.2863
I(MED_a_perc - 68.72)	-0.0229	0.0627	-0.3648	0.7153	-0.1457	0.1000
I(MED_b_perc - 8.536)	0.1342	0.1142	1.1748	0.2401	-0.0897	0.3581
I(MED_t_perc - 11.1)	-0.0591	0.0758	-0.7793	0.4358	-0.2078	0.0895

N.B.: predictor abbreviations refer to percentage of patients treated with atypical (MED_a_perc) or typical (MED_t_perc) antipsychotics, both antipsychotic medication types (MED_b_perc) and the percentage of unmedicated patients (intrcpt, i.e. baseline).

- n) Left MOFC thickness differences by antipsychotic medication type percentages, covarying for gender and age

We also investigated whether left MOFC thickness differed by the percentage of patients treated with typical or atypical antipsychotics, both antipsychotic medication types and the percentage of unmedicated patients, when additionally controlling for gender and age. Shown below are the estimate, the standard error, p value and the confidence intervals of mean left MOFC thickness for each antipsychotic medication group. Percentages were mean centered and compared to unmedicated patients (baseline). Results were based on estimates from 13 studies.

Mixed-Effects Model (k = 13; tau ² estimator: REML)	
tau ² (estimated amount of residual heterogeneity):	0.0076 (SE = 0.0043)
tau (square root of estimated tau ² value):	0.0873
I ² (residual heterogeneity / unaccounted variability):	96.44%
H ² (unaccounted variability / sampling variability):	28.06
R ² (amount of heterogeneity accounted for):	30.46%

Test for Residual Heterogeneity:

QE(df = 7) = 235.2282, p-val < .0001

Test of Moderators (coefficient(s) 2,3,4,5,6):

QM(df = 5) = 10.1896, p-val = 0.0700

Model Results:

	estimate	se	zval	pval	ci.lb	ci.ub
intrcpt ***	2.4799	0.0267	92.8312	<.0001	2.4276	2.5323
I(Age_mean - 34.72)	-0.0041	0.0100	-0.4115	0.6807	-0.0237	0.0155
I(MED_a_perc - 68.72) *	-0.0056	0.0022	-2.5524	0.0107	-0.0098	-0.0013
I(MED_b_perc - 8.536)	-0.0016	0.0046	-0.3361	0.7368	-0.0106	0.0075
I(MED_t_perc - 11.1)	-0.0053	0.0035	-1.5391	0.1238	-0.0121	0.0015
I(MF_ratio - 2.255)	-0.0091	0.0386	-0.2365	0.8130	-0.0848	0.0666

N.B.: predictor abbreviations refer to percentage of patients treated with atypical (MED_a_perc) or typical (MED_t_perc) antipsychotics, both antipsychotic medication types (MED_b_perc) and the percentage of unmedicated patients (intrcpt, i.e. baseline) as well as the ratio of males versus females (MF_ratio).

- o) Left MOFC thickness differences by antipsychotic medication type percentages, covarying for gender and length of illness

We also investigated whether left MOFC thickness differed by the percentage of patients treated with typical or atypical antipsychotics, both antipsychotic medication types and the percentage of unmedicated patients, when additionally controlling for gender and length of illness. Shown below are the estimate, the standard error, p value

and the confidence intervals of mean left MOFC thickness for each antipsychotic medication group. Percentages were mean centered and compared to unmedicated patients (baseline). Results were based on estimates from 11 studies.

Mixed-Effects Model (k = 11; tau^2 estimator: REML)

tau^2 (estimated amount of residual heterogeneity):

0.0087 (SE = 0.0057)

tau (square root of estimated tau^2 value):

0.0932

I^2 (residual heterogeneity / unaccounted variability):

97.07%

H^2 (unaccounted variability / sampling variability):

34.16

R^2 (amount of heterogeneity accounted for):

31.69%

Test for Residual Heterogeneity:

QE(df = 5) = 187.9049, p-val < .0001

Test of Moderators (coefficient(s) 2,3,4,5,6):

QM(df = 5) = 9.6171, p-val = 0.0868

Model Results:

	estimate	se	zval	pval	ci.lb	ci.ub
intrcpt	2.4693	0.0348	70.9461	<.0001	2.4011	2.5376

I(LOI_mean - 10.74)	-0.0077	0.0119	-0.6514	0.5148	-0.0310	0.0155
I(MED_a_perc - 68.72)	-0.0051	0.0025	-2.0599	0.0394	-0.0100	-0.0002
*						
I(MED_b_perc - 8.536)	0.0013	0.0074	0.1793	0.8577	-0.0132	0.0159
I(MED_t_perc - 11.1)	-0.0042	0.0036	-1.1451	0.2522	-0.0113	0.0030
I(MF_ratio - 2.255)	0.0089	0.0644	0.1384	0.8899	-0.1172	0.1350

N.B.: predictor abbreviations refer to percentage of patients treated with atypical (MED_a_perc) or typical (MED_t_perc) antipsychotics, both antipsychotic medication types (MED_b_perc) and the

percentage of unmedicated patients (intrcpt, i.e. baseline) as well as the ratio of males versus females (MF_ratio) and mean length of illness (LOI_mean).

p) Moderation analysis of single vs multisite status on the association between global SANS scores and left MOFC thickness

Shown below are the estimate, the standard error, p value and the confidence intervals for the moderating effects of single vs multisite status on the meta-analytical standardized regression beta of global SANS scores on left MOFC thickness. The intercept refers to the effect of global SANS score on left MOFC thickness after accounting for single vs multisite status. Results were based on estimates from 17 studies.

Mixed-Effects Model (k = 17; tau² estimator: REML)

tau² (estimated amount of residual heterogeneity): 0.0078 (SE = 0.0066)

tau (square root of estimated tau² value): 0.0886

I² (residual heterogeneity / unaccounted variability): 45.43%

H² (unaccounted variability / sampling variability): 1.83

R² (amount of heterogeneity accounted for): 0.00%

Test for Residual Heterogeneity:

QE(df = 15) = 25.9669, p-val = 0.0384

Test of Moderators (coefficient(s) 2):

QM(df = 1) = 0.6443, p-val = 0.4222

Model Results:

	estimate	se	zval	pval	ci.lb	ci.ub
intrcpt *	-0.0954	0.0409	-2.3295	0.0198	-0.1756	-0.0151
as.factor(multisite)1	0.0567	0.0706	0.8027	0.4222	-0.0817	0.1950

q) Effects of covariates: left MOFC <- global SANS + CPZ + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of current chlorpromazine equivalents (CPZ, based on Woods *et al.* (2003)) on left MOFC thickness. Results were based on estimates from 12 studies.

Random-Effects Model (k = 12; tau^2 estimator: REML)					
tau^2 (estimated amount of total heterogeneity): 0.0003 (SE = 0.0040)					
tau (square root of estimated tau^2 value): 0.0160					
I^2 (total heterogeneity / total variability): 2.26%					
H^2 (total variability / sampling variability): 1.02					
Test for Heterogeneity:					
Q(df = 11) = 9.7194, p-val = 0.5558					
Model Results:					
estimate	se	zval	pval	ci.lb	ci.ub
-0.0413	0.0301	-1.3717	0.1701	-0.1003	0.0177

r) Effects of covariates: left MOFC <- global SANS + CPZ + gender + age + site (if applicable)

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of global SANS scores on left MOFC thickness, controlling for current chlorpromazine equivalents (based on Woods *et al.* (2003)). Results were based on estimates from 12 studies.

Random-Effects Model (k = 12; tau ² estimator: REML)						
tau ² (estimated amount of total heterogeneity): 0.0120 (SE = 0.0104)						
tau (square root of estimated tau ² value): 0.1093						
I ² (total heterogeneity / total variability): 52.08%						
H ² (total variability / sampling variability): 2.09						
Test for Heterogeneity:						
Q(df = 11) = 22.4039, p-val = 0.0214						
Model Results:						
estimate	se	zval	pval	ci.lb	ci.ub	
-0.0950	0.0462	-2.0564	0.0397	-0.1856	-0.0045	*

- s) Effects in DSM-IV schizophrenia subtypes or schizoaffective / -phreniform disorder only:
left MOFC <- global SANS + CPZ + gender + age + site (if applicable)

Schizophrenia inclusion criteria were site-specific. In detail, five sites included only patients with DSM-IV subtypes of schizophrenia. Eight sites also included schizoaffective and schizopreniform

patients in their samples, while only a minority of sites ($N_{\text{sites}}=4$; 7% of our total sample) also included a few patients with other psychotic disorders, such as psychotic disorder NOS.

Of those sites where individual diagnostic codes for patients were available to us ($N_{\text{sites}}=11$, including the four sites with patients with other psychotic disorders), 77% of patients were diagnosed with Schizophrenia subtypes and 12% with schizoaffective / -phreniform disorders.

To ensure that our results were not influenced by including patients with other psychotic disorders, we repeated the analyses including only patients with DSM-IV schizophrenia subtypes or schizoaffective / -phreniform disorders.

Shown below are the estimate, the standard error, p value and the confidence intervals for the meta-analytical standardized regression beta of global SANS scores on left MOFC thickness, including only patients with DSM-IV schizophrenia subtypes or schizoaffective / -phreniform disorders. Results were based on estimates from 16 studies.

Random-Effects Model ($k = 16$; τ^2 estimator: REML)

τ^2 (estimated amount of total heterogeneity): 0.0060 (SE = 0.0061)

τ (square root of estimated τ^2 value): 0.0774

I^2 (total heterogeneity / total variability): 36.24%

H^2 (total variability / sampling variability): 1.57

Test for Heterogeneity:

$Q(df = 15) = 22.6004$, $p\text{-val} = 0.0930$

Model Results:

estimate	se	zval	pval	ci.lb	ci.ub	
-0.0780	0.0339	-2.2992	0.0215	-0.1445	-0.0115	*

SM 2.2 Exploratory analyses on ten more frontal regions

Univariate models were used to analyse the effect of negative symptoms on ten other frontal regions (separately for the left and right hemisphere; see SM Table 3s), controlling for sex, age and site (if applicable). After Bonferroni correction for multiple testing (ten regions per hemisphere = 20 tests), two regions remained significant on the left hemisphere, while no effect was observed in any of the regions on the right side (SM Table 3s).

region-of-interest	<i>left hemisphere</i>		<i>right hemisphere</i>		Nsites
	β std	<i>p</i>	β std	<i>p</i>	
caudalmiddlefrontal	-0.0196	0.5052	-0.0474	0.1406	17
lateralorbitofrontal	-0.0756	0.0017	-0.0732	0.0114	17
caudalanteriorcingulate	-0.0164	0.5169	-0.0547	0.0921	13
parsopercularis	-0.0821	0.001	-0.0357	0.2544	15
parsorbitalis	-0.0512	0.0512	-0.0479	0.0897	13
parstriangularis	-0.0659	0.0101	-0.0422	0.086	15
rostralanteriorcingulate	0.0137	0.6022	-0.0434	0.1884	13
rostralmiddlefrontal	-0.0509	0.0446	-0.0484	0.0613	17

Walton et al.		Negative symptoms and cortical thinning			
superiorfrontal	-0.0662	0.0223	-0.0640	0.0264	17
frontalpole	-0.0398	0.1149	0.0006	0.9821	13

SM Table 3s. Exploratory analyses on ten more frontal regions.

References

Woods, SW (2003). Chlorpromazine equivalent doses for the newer atypical antipsychotics. *The Journal of Clinical Psychiatry* **64**(6), 663–667.

SM 3 Acknowledgements

Support was provided by R01MH107703 and K23MH098130 to TDS, R01MH101111 to DHW, European Community's 7th Framework Programme (#602450, IMAGEMEND) to NTD. The CLING study and the HMS studies were partially supported by a research grant from the Competence Network Schizophrenia to Oliver Gruber. The Dublin study was supported by grant funding from the Irish Health Research Board (grant number HRA_POR/2012/54) and Science Foundation Ireland (grant numbers 12/IP/1359 and 08/IN.1/B1916). The Edinburgh study was supported by the Medical Research Council through a Clinical Training Fellowship (Ref G84/5699). Further funding was also provided by the Mortimer and Theresa Sackler Foundation. The FBIRN study was supported by the National Center for Research Resources at the National Institutes of Health (grant numbers: NIH 1 U24 RR021992 (Function Biomedical Informatics Research Network) and NIH 1 U24 RR025736-01 (Biomedical Informatics Research Network Coordinating Center; <http://www.birncommunity.org>). FBIRN data was processed by the UCI High Performance Computing cluster supported by Joseph Farran, Harry Mangalam, and Adam Brenner and the National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health, through Grant UL1 TR001414 UL1 TR001414. FBIRN thanks Mrs. Liv McMillan for overall study coordination. The NUI Galway study was supported by grant funding from the Health Research Board (grant number HRA_POR/2011/100) and the Wellcome Trust (grant number 072894/2/03/Z). The HUBIN study was supported by the Swedish Research Council (grant numbers K2009-62X-15077-06-3 and K2012-61X-15077-09-3), the Karolinska Institutet and the Knut and Alice Wallenberg Foundation. The MCIC study was supported by the National Institutes of Health (NIH/NCRR P41RR14075 and R01EB005846 (to Vince D. Calhoun)), the Department of Energy (DE-FG02-99ER62764), the Mind Research Network, the Morphometry BIRN (1U24, RR021382A), the Function BIRN (U24RR021992-01, NIH.NCRR MO1 RR025758-01, NIMH 1RC1MH089257 to Vince D. Calhoun), the Deutsche Forschungsgemeinschaft (research fellowship to Esther Walton and Stefan Ehrlich), and a NARSAD Young Investigator Award (to Stefan Ehrlich). The PAFIP study was supported by Instituto de Salud Carlos III, FIS 00/3095, 01/3129, PI020499, PI060507, PI10/00183, the SENY Fundació Research Grant CI 2005-0308007, the Fundación Marqués de Valdecilla API07/011 and the IDIVAL Neuroimaging Unit in the technical execution of this work. The TOP study was supported by the Research Council of Norway (#213837, #217776, #223273), the South-East Norway Health Authority

(2013-123), and the KG Jebsen Foundation. The UMCU study was supported by the Dutch Organization for Health Research and Development ZonMW grants 90802123 and 91746370 (to Hilleke E. Hulshoff Pol) and 10-000-1001 (to René S. Kahn) and NWO/ ZonMW Innovation Impulse (VIDI) nr. 017.106.301 (Iris Sommer) . The UPENN study was supported by National Institute of Mental Health grants MH064045, MH60722, MH019112, and MH085096 (DHW). Theodore D. Satterthwaite was supported by MH098130 and by the Marc Rapport Family through NARSAD. Research reported in this publication was supported by the National Institute of Biomedical Imaging and Bioengineering (NIBIB) of the National Institutes of Health under Award Number U54EB020403. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Osaka study was partially supported by Brain/MINDS and Health and Labour Sciences Research Grants for Comprehensive Research on Persons with Disabilities from AMED and KAKENHI from MEXT. The IRCCS Santa Lucia Foundation of Rome Study was partially supported by the Italian Ministry of Health (RC12-13-14-15A Grant) and by the by the European Commission ERA-Net NEURON joint transnational call 2010 (European Research Projects on Mental Disorders: NEUCONNECT). The KaSP consortium was supported by grants from: the Swedish Research Council (grant nr. 2011-4730 and samverkansanslag), the Stockholm County Council (grant nr. ALF 20090192, ALF 20140469, ALF 20150475) and the Centre for Psychiatric Research (grant nr. CPF 100/2011) Swedish Medical Research Council (2009-7053; 2013-2838), the Swedish Brain foundation, Söderström Königska and Torsten Söderbergs Stiftelse.